


# ENSO: Recent Evolution, Current Status and Predictions



Update prepared by:  
Climate Prediction Center / NCEP  
29 December 2014

# Outline

Summary

Recent Evolution and Current Conditions

Oceanic Niño Index (ONI)

Pacific SST Outlook

U.S. Seasonal Precipitation and Temperature Outlooks

Summary

# Summary

## ENSO Alert System Status: El Niño Watch

ENSO-neutral conditions continue.\*

Positive equatorial sea surface temperature (SST) anomalies continue across most of the Pacific Ocean.

There is an approximately 65% chance that El Niño conditions will be present during the Northern Hemisphere winter and last into the Northern Hemisphere spring 2015.\*


\* Note: These statements are updated once a month in association with the ENSO Diagnostics Discussion, which can be found by clicking [here](#).

# Recent Evolution of Equatorial Pacific SST Departures (°C)

From January- February 2014, SSTs were mostly below average across the eastern equatorial Pacific.

From March-June 2014, above-average SSTs (departures  $>0.5^{\circ}\text{C}$ ) were evident near the Date Line and in the eastern Pacific.


Recently, positive SST anomalies cover most of the equatorial Pacific.



# Niño Region SST Departures (°C) Recent Evolution


The latest weekly SST departures are:

Niño 4	0.9°C
Niño 3.4	0.7°C
Niño 3	0.7°C
Niño 1+2	-0.2°C




# SST Departures (°C) in the Tropical Pacific During the Last Four Weeks

During the last four weeks, equatorial SSTs were above average across most of the Pacific.



# Global SST Departures (°C) During the Last Four Weeks


During the last four weeks, equatorial SSTs were above average across most of the Pacific, and below average in the Atlantic Ocean.



# Weekly SST Departures during the Last Four Weeks

During the last four weeks, equatorial SST anomalies remained positive over most of the Pacific, except near zero near South America.


## Weekly SST Anomalies (DEG C)





# Change in Weekly SST Departures over the Last Four Weeks

During the last four weeks, changes in equatorial SST anomalies were negative across most of the eastern half of the Pacific.




# Upper-Ocean Conditions in the Equatorial Pacific

The basin-wide equatorial upper ocean (0-300 m) heat content is greatest prior to and during the early stages of a Pacific warm (El Niño) episode (compare top 2 panels), and least prior to and during the early stages of a cold (La Niña) episode.

The slope of the oceanic thermocline is least (greatest) during warm (cold) episodes.


Recent values of the upper-ocean heat anomalies (slightly positive) and thermocline slope index (near zero) reflect a tendency toward El Niño conditions.

*The monthly thermocline slope index represents the difference in anomalous depth of the 20°C isotherm between the western Pacific (160°E-150°W) and the eastern Pacific (90°-140°W).*




# Central and Eastern Pacific Upper-Ocean (0-300 m) Weekly Average Temperature Anomalies

Subsurface temperature anomalies strongly increased during January - March 2014 and then decreased. Temperature anomalies then increased from late July to late August, were relatively unchanged from September to mid-October, and increased from mid-October to mid-November. Recently, the anomalies have decreased.




# Sub-Surface Temperature Departures in the Equatorial Pacific

Since late October, positive subsurface temperature anomalies have stretched across most of the equatorial Pacific.



During December, positive subsurface anomalies in the central Pacific weakened.




# Tropical OLR and Wind Anomalies During the Last 30 Days

Positive OLR anomalies (suppressed convection and precipitation, red shading) were apparent near the Date Line (centered south of the equator). Negative OLR anomalies (enhanced convection and precipitation) were evident over Indonesia, the Philippines, and northern Australia.

Low-level (850-hPa) winds were near-average across most of the equatorial Pacific, while anomalous easterlies were present near Papua New Guinea.

Anomalous upper-level (200-hPa) easterly winds were evident across the east-central equatorial Pacific.



# Intraseasonal Variability

Intraseasonal variability in the atmosphere (wind and pressure), which is often related to the Madden-Julian Oscillation (MJO), can significantly impact surface and subsurface conditions across the Pacific Ocean.

Related to this activity:

Significant weakening of the low-level easterly winds usually initiates an eastward-propagating oceanic Kelvin wave.

# Weekly Heat Content Evolution in the Equatorial Pacific


During January - May 2014, the downwelling phase of a strong Kelvin wave crossed the Pacific.

During May-July, positive temperature anomalies progressively disappeared from the equatorial Pacific in response to the upwelling phase of the Kelvin wave.

During October-November, positive subsurface temperature anomalies increased and shifted eastward in association with the downwelling phase of a Kelvin wave.

Since November, a weak upwelling phase of a Kelvin wave has shifted eastward.

Oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Down-welling and warming occur in the leading portion of a Kelvin wave, and up-welling and cooling occur in the trailing portion.






# Low-level (850-hPa) Zonal (east-west) Wind Anomalies ( $\text{m s}^{-1}$ )

Since mid November, easterly anomalies have persisted over the western equatorial Pacific.

Near the Date Line, weak westerly wind anomalies have been evident.

Westerly Wind Anomalies (orange/red shading)  
Easterly Wind Anomalies (blue shading)






# Upper-level (200-hPa) Velocity Potential Anomalies

During mid-August to mid-September and October-mid December, eastward propagating velocity potential anomalies were observed.

During November through early December, the Madden-Julian Oscillation (MJO) emerged.

Unfavorable for precipitation (brown shading)  
Favorable for precipitation (green shading)

200-hPa Velocity Potential Anomaly: 5N-5S  
5-day Running Mean




# Outgoing Longwave Radiation (OLR) Anomalies

During late July-early August 2014, positive OLR anomalies shifted eastward from the Indian Ocean, across Indonesia to near the Date Line.

Since late October, positive anomalies have been observed near the Date Line and over the western Pacific.

Drier-than-average Conditions (orange/red shading)  
Wetter-than-average Conditions (blue shading)



# Oceanic Niño Index (ONI)

The ONI is based on SST departures from average in the Niño 3.4 region, and is a principal measure for monitoring, assessing, and predicting ENSO.

Defined as the three-month running-mean SST departures in the Niño 3.4 region. Departures are based on a set of improved homogeneous historical SST analyses (Extended Reconstructed SST - ERSST.v3b). The SST reconstruction methodology is described in Smith et al., 2008, J. Climate, vol. 21, 2283-2296.)

Used to place current events into a historical perspective

NOAA's operational definitions of El Niño and La Niña are keyed to the ONI index.

# NOAA Operational Definitions for El Niño and La Niña

El Niño: characterized by a positive ONI greater than or equal to  $+0.5^{\circ}\text{C}$ .


La Niña: characterized by a negative ONI less than or equal to  $-0.5^{\circ}\text{C}$ .

By historical standards, to be classified as a full-fledged El Niño or La Niña episode, these thresholds must be exceeded for a period of at least 5 consecutive overlapping 3-month seasons.

CPC considers El Niño or La Niña conditions to occur when the monthly Niño3.4 OISST departures meet or exceed  $\pm 0.5^{\circ}\text{C}$  along with consistent atmospheric features. These anomalies must also be forecasted to persist for 3 consecutive months.

# ONI (°C): Evolution since 1950

The most recent ONI value (September - November 2014) is 0.5°C.



# Historical El Niño and La Niña Episodes Based on the ONI computed using ERSST.v3b

## El Niño

### Highest ONI Value

JJA 1951 - DJF 1951/52	1.2
DJF 1952/53 - JFM 1954	0.8
MAM 1957 - JJA 1958	1.8
OND 1958 - FMA 1959	0.6
MJJ 1963 - JFM 1964	1.4
AMJ 1965 - MAM 1966	1.9
JAS 1968 - DJF 1969/70	1.1
AMJ 1972 - FMA 1973	2.1
ASO 1976 - JFM 1977	0.8
ASO 1977 - JFM 1978	0.8
AMJ 1982 - MJJ 1983	2.2
JAS 1986 - JFM 1988	1.6
AMJ 1991 - MJJ 1992	1.6
ASO 1994 - FMA 1995	1.2
AMJ 1997 - MAM 1998	2.4
AMJ 2002 - JFM 2003	1.3
JJA 2004 - DJF 2004/05	0.7
ASO 2006 - DJF 2006/07	1.0
JJA 2009 - MAM 2010	1.6

## La Niña

### Lowest ONI Value

ASO 1949 - JAS 1950	-1.4
SON 1950 - JFM 1951	-0.8
AMJ 1954 - NDJ 1956/57	-1.7
AMJ 1964 - DJF 1964/65	-0.8
JJA 1970 - DJF 1971/72	-1.3
AMJ 1973 - JJA 1974	-2.0
SON 1974 - MAM 1976	-1.7
ASO 1983 - DJF 1983/84	-0.9
SON 1984 - ASO 1985	-1.1
AMJ 1988 - AMJ 1989	-1.9
ASO 1995 - FMA 1996	-0.9
JJA 1998 - FMA 2001	-1.7
OND 2005 - FMA 2006	-0.9
JAS 2007 - MJJ 2008	-1.5
OND 2008 - FMA 2009	-0.8
JJA 2010 - MAM 2011	-1.5
ASO 2011 - FMA 2012	-1.0

NOTE (Mar. 2012): The historical values of the ONI have slightly changed due to an update in the climatology. Please click [here](#) for more details on the methodology.

# Historical El Niño and La Niña Episodes Based on the ONI computed using ERSST.v3b


Recent Pacific warm (red) and cold (blue) episodes based on a threshold of +/- 0.5 °C for the Oceanic Niño Index (ONI) [3 month running mean of ERSST.v3b SST anomalies in the Niño 3.4 region (5N-5S, 120-170W)]. For historical purposes El Niño and La Niña episodes are defined when the threshold is met for a minimum of 5 consecutive over-lapping seasons. The complete table going back to DJF 1950 can be found [here](#).

Year	DJF	JFM	FMA	MAM	AMJ	MJJ	JJA	JAS	ASO	SON	OND	NDJ
2002	-0.2	0.0	0.1	0.3	0.5	0.7	0.8	0.8	0.9	1.2	1.3	1.3
2003	1.1	0.8	0.4	0.0	-0.2	-0.1	0.2	0.4	0.4	0.4	0.4	0.3
2004	0.3	0.2	0.1	0.1	0.2	0.3	0.5	0.7	0.8	0.7	0.7	0.7
2005	0.6	0.4	0.3	0.3	0.3	0.3	0.2	0.1	0.0	-0.2	-0.5	-0.8
2006	-0.9	-0.7	-0.5	-0.3	0.0	0.1	0.2	0.3	0.5	0.8	1.0	1.0
2007	0.7	0.3	-0.1	-0.2	-0.3	-0.3	-0.4	-0.6	-0.8	-1.1	-1.2	-1.4
2008	-1.5	-1.5	-1.2	-0.9	-0.7	-0.5	-0.3	-0.2	-0.1	-0.2	-0.5	-0.7
2009	-0.8	-0.7	-0.5	-0.2	0.2	0.4	0.5	0.6	0.8	1.1	1.4	1.6
2010	1.6	1.3	1.0	0.6	0.1	-0.4	-0.9	-1.2	-1.4	-1.5	-1.5	-1.5
2011	-1.4	-1.2	-0.9	-0.6	-0.3	-0.2	-0.2	-0.4	-0.6	-0.8	-1.0	-1.0
2012	-0.9	-0.6	-0.5	-0.3	-0.2	0.0	0.1	0.4	0.5	0.6	0.2	-0.3
2013	-0.6	-0.6	-0.4	-0.2	-0.2	-0.3	-0.3	-0.3	-0.3	-0.2	-0.3	-0.4
2014	-0.6	-0.6	-0.5	-0.1	0.1	0.1	0.0	0.0	0.2	0.5		

# CPC/IRI Probabilistic ENSO Outlook

Updated: 4 December 2014

The chance of El Niño is approximately 65% during the Northern Hemisphere winter and decreases into spring/summer 2015.





# IRI/CPC Pacific Niño

## 3.4 SST Model Outlook

Most models favor El Niño (greater than or equal to  $+0.5^{\circ}\text{C}$ ) to form and persist through Northern Hemisphere spring 2015.






Figure provided by the International Research Institute (IRI) for Climate and Society (updated 16 December 2014).

# SST Outlook: NCEP CFS.v2 Forecast (PDF corrected)

Issued: 29 December 2014


The CFS.v2 ensemble mean (black dashed line) predicts El Niño or warm-neutral conditions into mid-2015.



# Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

During late October - mid December, the pattern generally featured an anomalous ridge over western N. America and an anomalous trough over eastern N. America. This pattern led to above-average temperatures in the West, and below average temperatures in the East.

Since mid December, an anomalous ridge and warmer-than-average temperatures have been apparent over Canada and the northern half of the U.S.






# Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

During late October - mid December, the pattern generally featured an anomalous ridge over western N. America and an anomalous trough over eastern N. America. This pattern led to above-average temperatures in the West, and below average temperatures in the East.


Since mid December, an anomalous ridge and warmer-than-average temperatures have been apparent over Canada and the northern half of the U.S.



# Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

During late October - mid December, the pattern generally featured an anomalous ridge over western N. America and an anomalous trough over eastern N. America. This pattern led to above-average temperatures in the West, and below average temperatures in the East.


Since mid December, an anomalous ridge and warmer-than-average temperatures have been apparent over Canada and the northern half of the U.S.




# U.S. Temperature and Precipitation Departures During the Last 30 Days

End Date: 27 December 2014

### Percent of Average Precipitation




### Temperature Departures (degree C)




# U.S. Temperature and Precipitation Departures During the Last 90 Days

End Date: 27 December 2014

### Percent of Average Precipitation





### Temperature Departures (degree C)



# U. S. Seasonal Outlooks

## January - March 2015

The seasonal outlooks combine the effects of long-term trends, soil moisture, and, when appropriate, ENSO.





# Summary

## ENSO Alert System Status: El Niño Watch

ENSO-neutral conditions continue.\*

Positive equatorial sea surface temperature (SST) anomalies continue across most of the Pacific Ocean.

There is an approximately 65% chance that El Niño conditions will be present during the Northern Hemisphere winter and last into the Northern Hemisphere spring 2015.\*

\* Note: These statements are updated once a month in association with the ENSO Diagnostics Discussion, which can be found by clicking [here](#).